


Name:			
Enrolment No:			
UPES End Semester Examination, May 2023			
Course: Physical Chemistry IV Program: BSc (H) Chemistry & Int. B.Sc.-M.Sc. Chemistry Course Code: CHEM 2025		Semester: IV Time : 03 hrs. Max. Marks: 100	
Instructions: 1) Read all questions carefully. 2) Attempt all questions under one section in one place.			
SECTION A (5Qx4M= 20 Marks)			
S. No.		Marks	CO
Q 1	Calculate the value of an einstein of energy in electron volts for radiation of $3 \times 10^{13} \text{ s}^{-1}$ frequency.	4	CO1
Q2	At room temperature the molar conductance of H_2O is $1.0 \times 10^{-6} \text{ S m}^2 \text{ mol}^{-1}$ and the molar conductance at infinite dilution is $550 \text{ S m}^2 \text{ mol}^{-1}$. Calculate the degree of dissociation of water?	4	CO1
Q3	The rate constant for a reaction of zero order is $0.0030 \text{ mol L}^{-1} \text{ s}^{-1}$. How long will it take for the initial concentration to fall from 0.10 M to 0.075 M ?	4	CO1
Q4	Differentiate between (i) Specific conductance and Equivalent conductance (ii) Galvanic cell and Electrolytic cell	2 +2	CO1
Q5	Calculate equilibrium constant for the reaction $\text{Zn}_{(s)} + \text{Cu}^{2+}_{(aq)} \rightleftharpoons \text{Cu}_{(s)} + \text{Zn}^{2+}_{(aq)}$ at 25°C . Standard <i>emf</i> of the cell is 1.10 V .	4	CO1
SECTION B (4Qx10M= 40 Marks)			
Q6	Calculate the transmittance, absorbance and absorption coefficient of a solution which absorbs 90% of a certain wavelength of light beam passed through a 1 cm cell containing a 0.25 M solution.	10	CO2
Q7	Write a brief note on Norrish Type-I and Norrish Type-II reactions.	10	CO1
Q8	Ionization constant of acetic acid and ionic product of water at 25°C are 1.75×10^{-5} and 1×10^{-14} respectively. Calculate the hydrolysis constant of sodium acetate and its degree of hydrolysis in 0.1 molar solution at 25°C .	10	CO3
Q9	What is meant by transport number of an ion? How is it determined using Hittorf's method and Moving Boundary method?	10	CO2
<i>Or</i>			

	A first order gas reaction $A_2B_{2(g)} \rightarrow 2A_{(g)}$ at the temperature $400\text{ }^\circ\text{C}$ has the rate constant $k = 2.0 \times 10^{-4}\text{ s}^{-1}$. What percentage of A_2B_2 is decomposed on heating for 900 seconds?		
SECTION-C (2Qx20M=40 Marks)			
Q10	<p>(a) What do you understand by quantum yield of a photochemical reaction? Why do some reactions have high quantum yield whereas some others have very low value? What is the modified definition of Stark-Einstein law?</p> <p>(b) The resistance of a conductivity cell when filled with 0.02 M KCl solution is 164 ohms at 298 K. However, when filled with 0.05 M AgNO_3 solution, its resistance is found to be 78.5 ohms. If specific conductivity of 0.02 M KCl is $2.768 \times 10^{-3}\text{ ohm}^{-1}$, calculate</p> <p>(i) The conductivity of 0.05 M AgNO_3</p> <p>(ii) The molar conductivity of AgNO_3 solution</p>	10 +10	CO2
Q11	<p>(a) Prove that degree of hydrolysis of a salt of weak acid and weak base is independent of the concentration of the solution</p> <p style="text-align: center;"><i>Or</i></p> <p>Calculate the electrode potential of a copper wire dipped in 0.1 molar copper sulphate solution at $25\text{ }^\circ\text{C}$. At this temperature, the standard electrode potential of copper is 0.34 volt ($F = 96500\text{ coulombs}$; $R = 8.314\text{ J deg}^{-1}\text{ mol}^{-1}$). Assume CuSO_4 to be completely ionized and take the activity of copper ions equal to the molar concentration.</p> <p>(b) The decomposition of N_2O_5 to NO_2 and O_2 is first order with a rate constant of 4.8×10^{-4} per second at $45\text{ }^\circ\text{C}$.</p> <p>(i) if the initial concentration is $1.65 \times 10^{-2}\text{ mol/L}$, what is the concentration after 825 second?</p> <p>(ii) How long would it take for the concentration of N_2O_5 to decrease to $1.0 \times 10^{-2}\text{ mol/L}$ from its initial value, given in (i)?</p>	10 +10	CO3